arXiv:1703.10275v1 [math-ph] 30 Mar 2017

CONSTRUCTION OF A SET OF P-ADIC DISTRIBUTIONS

U. A. ROZIKOV, Z. T. TUGYONOV

Abstract. In this paper adapting to p-adic case some methods of real valued Gibbs measures on Cayley trees we construct several p-adic distributions on the set $\mathbb{Z}_p$ of p-adic integers. Moreover, we give conditions under which these p-adic distributions become p-adic measures (i.e. bounded distributions).

Mathematics Subject Classifications (2010). 46S10, 82B26, 12J12 (primary); 60K35 (secondary)

Key words. Cayley trees, p-adic numbers, p-adic distributions, p-adic measures.

1. Introduction

A p-adic distribution is an analogue of ordinary distributions that takes values in a ring of p-adic numbers [13]. Analogically to a measure on a measurable space, a p-adic measure is a special case of a p-adic distribution, i.e., a p-adic distribution taking values in a normed space is called a p-adic measure if the values on compact open subsets are bounded.

The main purpose for constructing p-adic measures is to integrate p-adic valued functions. The development parallels the classic treatment, beginning with the idea of Riemann sums. Note that such sums may not converge (even for continuous functions) if instead of p-adic measures one uses unbounded p-adic distributions. The theory of p-adic measures also useful in the context of p-adic L-functions following the works of B. Mazur (see [13] and [6] for details).

One of the first applications of p-adic numbers in quantum physics appeared in the framework of quantum logic [2]. Note that this model cannot be described by using conventional real-valued probability measures. In (real valued probability) measure theory Kolmogorov's extension theorem (see, e.g., [23, Ch. II]) is very helpful to construct measures.

A non-Archimedean analogue of the Kolmogorov's theorem was proved in [5]. Such a result allows to construct wide classes of p-adic distributions, stochastic processes and the possibility to develop statistical mechanics in the context of p-adic theory.

We refer the reader to [4], [8], [11], [10]- [19] where various p-adic models of statistical physics and p-adic distributions are studied.

In the present paper we construct several p-adic distributions and measures on the set $\mathbb{Z}_p$ of p-adic integers. We use some arguments of construction of real valued Gibbs measures on Cayley trees. To do this we present each element of the set $\mathbb{Z}_p$ by a path in the half (semi-infinite) Cayley tree. Then using known distribution relation (see equation (2.2)

below) we show that each collection of given p-adic distributions can be used to construct new distributions. Moreover, we give conditions under which these p-adic distributions become p-adic measures.

2. Preliminaries

2.1. **p-adic numbers.** Let $\mathbb{Q}$ be the field of rational numbers. For a fixed prime number p, every rational number $x \neq 0$ can be represented in the form $x = p^r \frac{n}{m}$, where $r, n \in \mathbb{Z}$, m is a positive integer, and n and m are relatively prime with p: $(p, n) = 1$, $(p, m) = 1$. The p-adic norm of x is given by

$$|x|_p = \begin{cases} p^{-r} & \text{for } x \neq 0 \\ 0 & \text{for } x = 0. \end{cases}$$

This norm is non-Archimedean and satisfies the so called strong triangle inequality

$$|x + y|_p \leq \max\{|x|_p, |y|_p\}.$$

The completion of $\mathbb{Q}$ with respect to the p-adic norm defines the p-adic field $\mathbb{Q}_p$. Any p-adic number $x \neq 0$ can be uniquely represented in the canonical form

$$x = p^{\gamma(x)}(x_0 + x_1 p + x_2 p^2 + \dots), \tag{2.1}$$

where $\gamma(x) \in \mathbb{Z}$ and the integers x_j satisfy: $x_0 > 0$, $0 \leq x_j \leq p - 1$. In this case $|x|_p = p^{-\gamma(x)}$.

The elements of the set $\mathbb{Z}_p = \{x \in \mathbb{Q}_p : |x|_p \leq 1\}$ are called p-adic integers.

We refer the reader to [13,22,27] for the basics of p-adic analysis and p-adic mathematical physics.

2.2. **p-adic distribution and measure.** Let $(X, \mathcal{B})$ be a measurable space, where $\mathcal{B}$ is an algebra of subsets of X. A function $\mu : \mathcal{B} \to \mathbb{Q}_p$ is said to be a p-adic *distribution* if for any $A_1, ..., A_n \in \mathcal{B}$ such that $A_i \cap A_j = \emptyset$, $i \neq j$, the following holds:

$$\mu(\bigcup_{j=1}^{n} A_j) = \sum_{j=1}^{n} \mu(A_j).$$

A p-adic distribution μ is called *measure* if it is bounded, i.e.

$$|\mu(A)|_p \leq +\infty, \quad \forall A \in \mathcal{B}.$$

A measure is called a probability measure if $\mu(X) = 1$, see, e.g. [9], [20].

In the metric space $\mathbb{Q}_p$ (with metric $\rho(x, y) = |x - y|_p$) basis of open sets can be taken as the following system of sets:

$$a + (p^N) = \left\{ x \in \mathbb{Q}_p : |x - a|_p \leq \frac{1}{p^N} \right\}, \quad a \in \mathbb{Q}_p, \quad N \in \mathbb{Z}.$$

Such a set is called an interval. The following theorem is known:

Theorem 1. *(see [13]) Every map μ of the set of intervals contained in $X \subset \mathbb{Q}_p$, for which*

$$\mu(a+(p^n)) = \sum_{b=0}^{p-1} \mu(a+bp^n+(p^{n+1})) \tag{2.2}$$

for any $a+(p^n) \subset X$, can be uniquely extended to a p-adic distribution on X.

Equation (2.2) is called a distribution relation.

Remark 1. *If $\mu_1, \mu_2, \dots, \mu_m$ are given distributions with values in $\mathbb{Q}_p$ and $\lambda_1, \lambda_2, \dots, \lambda_m \in \mathbb{Q}_p$ are arbitrary constants. Then*

$$\mu = \sum_{i=1}^{m} \lambda_i \mu_i \tag{2.3}$$

is a distribution. Indeed, define

$$\mu(a+(p^n)) = \sum_{i=1}^{m} \lambda_i \mu_i(a+(p^n))$$

then μ satisfies (2.2) since each μ_i satisfies (2.2). In the theory of real valued measures if one considers a convex set of such measures then to describe all points of this set it will be sufficient to know all extreme points of the set. In such a situations measures of the form (2.3) are not extreme and can be considered as linear combination of extreme measures. In the theory of (real) Gibbs measures for a given Hamiltonian (see [7] for details) the set of all such measures is a convex compact set. To construct new extreme points of this set in [3] the authors used some known extreme points. In this paper we will use idea of [3] to construct (see Theorem 2 below) new p-adic distributions from the known ones. Proof of Theorem 3 also reminds the construction of a limiting Gibbs measure with a given boundary condition on a Cayley tree (see [21] for details of the theory of Gibbs measures on trees).

2.3. **Cayley tree.** The Cayley tree (Bethe lattice [1]) Γ^k of order $k \geq 1$ is an infinite tree, i.e., a graph without cycles, such that exactly $k+1$ edges originate from each vertex. Let $\Gamma^k = (V, L)$ where V is the set of vertices and L the set of edges. Two vertices x and y are called *nearest neighbors* if there exists an edge $l \in L$ connecting them. We will use the notation $l = \langle x, y \rangle$. A collection of nearest neighbor pairs $\langle x, x_1 \rangle, \langle x_1, x_2 \rangle, ..., \langle x_{d-1}, y \rangle$ is called a *path* from x to y. The distance $d(x, y)$ on the Cayley tree is the number of edges of the shortest path from x to y.

If an arbitrary edge $\langle x^0, x^1 \rangle = l \in L$ is deleted from the Cayley tree Γ^k, it splits into two components– two semi-infinite trees Γ_0^k and Γ_1^k (each of which is called a half tree). In this paper we consider half tree $\Gamma_0^k = (V^0, L^0)$ (see Fig. 1).

For a fixed $x^0 \in V^0$, called the root, we set

$$W_n = \{x \in V^0 | d(x, x^0) = n\}, \qquad V_n^0 = \bigcup_{m=0}^{n} W_m.$$



FIGURE 1. Half Cayley tree of order three. The bold path π has representation by sequence $121\ldots$. On the left (resp. right) side of this path the distribution μ coincides with μ_1 (resp. μ_2).

On the tree Γ^k one can introduce a partial ordering, by saying that $y > x$ if there exists a path $x = x_0, x_1, ..., x_n = y$ from x to y that "goes upwards", i.e., such that $d(x_m, x^0) = d(x_{m-1}, x^0) + 1, m = 1, \ldots, n$. The set of vertices $V_x = \{y \in V^0 | y \geq x\}$ and the edges connecting them form the semi-infinite tree Γ^k_x "growing" from the vertex $x \in V^0$.

Let p be a prime number. In this paper we consider the half Cayley tree of order p. Take an arbitrary (finite or infinite) path $x^0 = x_0 < x_1 < x_2 < \ldots$ on Γ^p_0 starting from the point x^0. We can represent the path by a sequence $i_0 i_1 i_2 \ldots$, where $i_n = 0, 1, \ldots, p-1$. Namely, we label by $l_0(x), l_1(x), \ldots, l_{p-1}(x)$ the edges going upward from the vertex $x \in V^0$. Then the path $x^0 = x_0 < x_1 < x_2 < \ldots$ can be assigned a sequence $i_0 i_1 \ldots$ such that $\langle x_{n-1}, x_n \rangle = l_{i_{n-1}}(x_{n-1})$, $n = 1, 2, \ldots$; the sequence $i_0 i_1 i_2 \ldots$ unambiguously determines the path $x^0 = x_0 < x_1 < x_2 < \ldots$.

A finite path $x^0 = x_0 < x_1 < x_2 < \cdots < x_n = x$ of length n determines a point $x \in W_n$. If the path $x^0 = x_0 < x_1 < x_2 < \cdots < x_n = x$ is represented by the sequence $i_0 i_1 \ldots i_n$, we assume that the vertex x is also represented by this sequence (see Fig. 1).

Let $x, y \in W_n$, and let x be represented by the sequence $i_0 i_1 \ldots i_n$ and y by $j_0 j_1 \ldots j_n$, where $i_0 = j_0$, $i_1 = j_1, \ldots$, $i_m = j_m$, but $i_{m+1} < j_{m+1}$ for some m. In this case, we write $x \prec y$. Similarly, for pathes π_1 represented by a sequence $i_0 i_1 \ldots i_n \ldots$ and π_2 represented by $j_0 j_1 \ldots j_n \ldots$, we write $\pi_1 \prec \pi_2$ if $i_0 = j_0$, $i_1 = j_1, \ldots$, $i_m = j_m$, but $i_{m+1} < j_{m+1}$ for some m.

Let $\pi = \{x^0 = x_0 < x_1 < \ldots\}$ be an infinite path, represented by the sequence $i_0 i_1 i_2 \ldots$. We assign the p-adic number

$$t = t(\pi) = \sum_{n=0}^{\infty} i_n p^n, \quad t \in \mathbb{Z}_p$$

to the path π. This assignment is a 1-1 correspondence. Moreover, the partial order $\prec$ on the set of paths can be used to obtain partial order on $\mathbb{Z}_p$: namely, for $t_1, t_2 \in \mathbb{Z}_p$ we write $t_1 \prec t_2$ iff they correspond to π_1,π_2 with $\pi_1 \prec \pi_2$.

3. Construction of distributions

Several examples of p-adic distributions are known: Haar's, Mazur's, Bernoulli's distributions etc. [13]. Recently some periodic distributions were constructed in [24]- [26].

Here for any two p-adic distribution given on $\mathbb{Z}_p$, say μ_1 and μ_2, $\mu_1 \neq \mu_2$ we construct a new distribution $\mu = \mu(\mu_1, \mu_2)$ defined on $\mathbb{Z}_p$.

Fix a path π with representation $\pi = i_0 i_1 \dots$. Take $a \in \mathbb{Z}_p$. If there exists $m \geq 0$ such that a has representation $a = i_0 + i_1 p + \dots + i_m p^m + a_{m+1} p^{m+1} + \dots$ with $a_{m+1} \neq i_{m+1}$, then we denote this m by $m(a, \pi)$. We say that two pathes starting from x^0 has non-empty intersection if they have at least one common edge.

Theorem 2. *Let $\pi = \{x^0 = x_0 < x_1 < \dots\}$ be an infinite path, represented by the sequence $i_0 i_1 i_2 \dots$ and μ_1, μ_2 are given distributions on $\mathbb{Z}_p$ such that $\mu_1 \neq \mu_2$,*

$$\mu_1(a + (p^n)) = \mu_2(a + (p^n)), \forall n \leq m(a, \pi) + 1, \tag{3.1}$$

then there is a distribution μ on $\mathbb{Z}_p$, which on the intervals is defined as follows

$$\mu(a + (p^n)) = \begin{cases} \mu_1(a + (p^n)), & \text{if } \pi_a \prec \pi, \pi_a \cap \pi = \emptyset \text{ or } \pi_a \cap \pi \neq \emptyset, \quad n \leq m \\ & \quad \text{or } \pi_a \cap \pi \neq \emptyset, \quad n \geq m+1,\ a_{m+1} < i_{m+1} \\ \mu_2(a + (p^n)), & \text{if } \pi \prec \pi_a, \pi_a \cap \pi = \emptyset \text{ or} \\ & \quad \pi_a \cap \pi \neq \emptyset, \quad n \geq m+1,\ a_{m+1} > i_{m+1} \end{cases} \tag{3.2}$$

where $m = m(a, \pi)$ and π_a is the path representing $a \in \mathbb{Z}_p$ (see Fig. 1).

Proof. Consider the following (all possible) cases:

Case I: $\pi_a \cap \pi = \emptyset$. In this case the path π_a always remains on the "left side" (i.e. $a_0 < i_0$) or always on the "right side" (i.e. $a_0 > i_0$) from the path π. For any $n \geq 1$ and any $b = 0, 1, \dots, p-1$ the number bp^n (present in formula (2.2)) makes contributions only on levels n, $n+1$, ... of the branch V_{a_0} growing from the vertex of the edge numbered by a_0, (where $a_0 \neq i_0$) of the tree Γ_0^p. We note that V_{a_0} does not intersect with π. Consequently we have $\pi_{a+bp^n} \cap \pi = \emptyset$. By definition of μ given in (3.2) in case $\pi_a \cap \pi = \emptyset$ and $\pi_{a+bp^n} \cap \pi = \emptyset$ for any $b = 0, 1, ..., p-1$ one easily checks that μ always coincides with μ_1 (or μ_2), hence it satisfies the distribution relation (2.2).

In case $n = 0$, the exists $b_0 \in \{0, 1, \dots, p-1\}$ such that $\pi_{a+b_0} \cap \pi \neq \emptyset$, i.e. $a + b_0 = i_0 + a_1 p + ...$, where i_0 is the first number in the presentation of π. In this case assuming $\pi_a \prec \pi$ the equality (2.2) can be written as

$$\mu_1(a + (p^0)) = \sum_{b=0}^{b_0} \mu_1(a + b + (p^1)) + \sum_{b=b_0+1}^{p-1} \mu_2(a + b + (p^1)). \tag{3.3}$$

Since the equality (2.2) is true for μ_1, the equality (3.3) is satisfied iff

$$\sum_{b=b_0+1}^{p-1}\mu_2(a+b+(p^1))=\sum_{b=b_0+1}^{p-1}\mu_1(a+b+(p^1)). \tag{3.4}$$

The last equality is true by condition (3.1).

Let now $\pi\prec\pi_a$ then the equality (2.2) can be written as

$$\mu_2(a+(p^0))=\sum_{b=0}^{b_0}\mu_2(a+b+(p^1))+\sum_{b=b_0+1}^{p-1}\mu_1(a+b+(p^1)). \tag{3.5}$$

Similarly as the case $\pi_a\prec\pi$ we obtain condition (3.4).

Case II: $\pi_a\cap\pi\neq\emptyset$. In this case there exists $m=m(a,\pi)\geq 0$ such that a has representation $a=i_0+i_1p+\cdots+i_mp^m+a_{m+1}p^{m+1}+\dots$ with $a_{m+1}\neq i_{m+1}$.

Subcase $n\geq m+2$: Since the path $a_{m+1}a_{m+2}\dots$ does not intersect with π, the equation (2.2) for $n\geq m+2$ is

$$\mu_1(i_0+i_1p+\cdots+i_mp^m+a_{m+1}p^{m+1}+\cdots+(p^n))=$$

$$\sum_{b=0}^{p-1}\mu_1(i_0+i_1p+\cdots+i_mp^m+a_{m+1}p^{m+1}+\cdots+bp^n+(p^{n+1})),\quad\text{if}\quad a_{m+1}<i_{m+1};$$

and

$$\mu_2(i_0+i_1p+\cdots+i_mp^m+a_{m+1}p^{m+1}+\cdots+(p^n))=$$

$$\sum_{b=0}^{p-1}\mu_2(i_0+i_1p+\cdots+i_mp^m+a_{m+1}p^{m+1}+\cdots+bp^n+(p^{n+1})),\quad\text{if}\quad a_{m+1}>i_{m+1}.$$

Since both μ_1 and μ_2 are distributions the last two equalities are true.

Subcase $n\leq m+1$: In this case by our condition (3.1) we have $\mu_1=\mu_2$, consequently, the distribution relation (2.2) is satisfied. □

The following theorem says that one can construct a new distribution using a collection of given distributions.

Theorem 3. *Let k be a natural number. Assume for each $t\in W_k$ a distribution ν_t on $\mathbb{Z}_p$ is given and there exists an interval $c+(p^{n_0})$, $n_0\geq k$ such that $\nu_t(c+(p^{n_0}))\neq\nu_s(c+(p^{n_0}))$ for at least one pair $t\neq s$, $t,s\in W_k$. Then there exists a distribution $\mu^{(k)}=\mu^{(k)}[\nu_t,t\in W_k]$ on $\mathbb{Z}_p$ which is different from ν_t for each $t\in W_k$.*

Proof. The distribution $\mu^{(k)}$ can be constructed as follows (see Fig. 2).

Case: $n\geq k$. Recall V_t is the set of vertices in the semi-infinite tree Γ_t^p growing from the vertex $t\in V^0$. In the case $n\geq k$ if $\pi_a\cap V_t\neq\emptyset$, for some $t\in W_k$ then π_{a+bp^n} also remains in V_t for any $b=0,1,\dots,p-1$. Therefore we define

$$\mu^{(k)}(a+(p^n))=\nu_t(a+(p^n)),\quad\text{if}\quad\pi_a\cap V_t\neq\emptyset, \tag{3.6}$$

then

$$\mu^{(k)}(a+bp^n+(p^{n+1}))=\nu_t(a+bp^n+(p^{n+1})),\quad\text{for each}\quad b=0,1,\dots,p-1.$$



FIGURE 2. The half Cayley tree of order three. Case $k = 1$, elements of W_1 are defined by $0, 1, 2$ and $\mu^{(1)}$ coincides with distribution ν_i, on the ith branch, where $i = 0, 1, 2$.

Since ν_t is a given distribution the equation (2.2) is satisfied.

Case: $n \le k-1$. Without loss of generality we define $\mu^{(k)}(a+(p^n))$ only for a which has the form

$$a = a_0 + a_1 p + \cdots + a_{k-1}p^{k-1},$$

where $a_i \in \{0, 1, \ldots, p-1\}$. Note that the sequence $a_0 a_1 \ldots a_{k-1}$ presents a unique point of W_k.

For $n = k-1$ define

$$\mu^{(k)}(a_0 + a_1 p + \cdots + a_{k-1}p^{k-1} + (p^{k-1}))$$
$$= \sum_{b=0}^{p-1} \nu_{a_0 a_1 \ldots a_{k-2}(a_{k-1}+b) \bmod p}(a_0 + a_1 p + \cdots + [(a_{k-1}+b) \bmod p]p^{k-1} + (p^k)). \quad (3.7)$$

Now using formula (3.7) we define (for $n = k-2$)

$$\mu^{(k)}(a_0 + a_1 p + \cdots + a_{k-2}p^{k-2} + (p^{k-2}))$$
$$= \sum_{b=0}^{p-1} \mu^{(k)}(a_0 + a_1 p + \cdots + [(a_{k-2}+b) \bmod p]p^{k-2} + (p^{k-1})) \quad (3.8)$$

Iterating this procedure we recurrently define

$$\mu^{(k)}(a_0 + a_1 p + \cdots + a_{k-m}p^{k-m} + (p^{k-m}))$$
$$= \sum_{b=0}^{p-1} \mu^{(k)}(a_0 + a_1 p + \cdots + [(a_{k-m}+b) \bmod p]p^{k-m} + (p^{k-m+1})), \quad (3.9)$$

where $m = 2, 3, \ldots, k$.

From the equalities (3.6)-(3.9) it follows that $\mu^{(k)}(a+(p^n))$ satisfies (2.2) for any $a \in \mathbb{Z}_p$ and any $n \ge 0$. Therefore, by Theorem 2 it follows that $\mu^{(k)}$ can be uniquely extended to a p-adic distribution on $\mathbb{Z}_p$.

Now we show that $\mu^{(k)}$ is different from ν_t for any $t \in W_k$. Let us assume the converse, i.e. there exists $u \in W_k$ such that $\mu^{(k)} = \nu_u$. This means that for any $a \in \mathbb{Z}_p$ and any $n \in \mathbb{N}$ we have

$$\mu^{(k)}(a + (p^n)) = \nu_u(a + (p^n)). \tag{3.10}$$

For any $a \in \mathbb{Z}_p$ there exists $t_a \in W_k$ such that $\pi_a \cap V_{t_a} \neq \emptyset$. Consequently, from (3.6) and (3.10) we get

$$\mu^{(k)}(a + (p^n)) = \nu_{t_a}(a + (p^n)) = \nu_u(a + (p^n)), \quad n \geq k. \tag{3.11}$$

When a runs arbitrary t_a takes all values on W_k. Therefore from (3.11) we get that $\nu_t(a + (p^n)) = \nu_u(a + (p^n))$, $n \geq k$ for any $t \in W_k$. But this is contradiction to the condition of theorem for $a = c$ and $n = n_0$. Thus it follows that $\mu^{(k)} \neq \nu_t$ for each $t \in W_k$. This completes the proof. □

Remark 2. *We note that the distribution $\mu^{(k)}$ mentioned in Theorem 3 is useful to construct two distributions μ_1 and μ_2 which satisfy the condition of Theorem 2. Indeed, let $k = 1$ and $\pi = i_0 i_1 \dots$, consider ν_0, ν_1 such that $\nu_0(c + (p^{n_0})) \neq \nu_1(c + (p^{n_0}))$, for some $n_0 \geq 1$ and $\pi_c \cap \pi = \emptyset$. Define $\tilde{\nu}_t$, $t \in W_1$ by*

$$\tilde{\nu}_t = \begin{cases} \nu_0, & \text{if } t = i_0 \\ \nu_1, & \text{if } t \neq i_0. \end{cases}$$

Take $\mu_1 = \nu_0$ and $\mu_2 = \mu^{(1)}[\tilde{\nu}_t, t \in W_1]$, it is easy to check that $\mu_1 \neq \mu_2$ and they satisfy the condition of Theorem 2.

In the p-adic integral theory one need to p-adic measure. Because integral of a continuous function calculated with respect to unbounded p-adic distribution may not exist (see [13] for details).

In the following theorem we give a conditions under which p-adic distributions mentioned in Theorems 2 and 3 become p-adic measures.

Theorem 4. *The following statements hold*

1) *Distribution μ, mentioned in Theorem 2, is bounded iff μ_1 and μ_2 are bounded.*
2) *Distribution $\mu^{(k)}$, mentioned in Theorem 3, is bounded iff $\nu_t(a + (p^n))$ is bounded for each $t \in W_k$ and $n \geq k$.*

Proof. 1) This is consequence of the equality (3.2).

2) Case $n \geq k$: In this case from (3.6) we get that $\mu^{(k)}$ is bounded iff ν_t is bounded for each $t \in W_k$.

Case $n \leq k-1$: Since the right hand side of (3.7) is sum of bounded terms (corresponding to the case $n = k$) the left hand side also should be bounded. Consequently, iterating of the recursions (3.7)-(3.9) one can see that $\mu^{(k)}$ is bounded iff $\nu_t(a + (p^n))$ is bounded for each $t \in W_k$ and $n \geq k$. □

Acknowledgements

This work was partially supported by Kazakhstan Ministry of Education and Science, grant 0828/GF4.

References

[1] Baxter R.J. *Exactly Solved Models in Statistical Mechanics* (Academic, London, 1982).
[2] Beltrametti E.G., Cassinelli G., Quantum Mechanics and p-adic Numbers,*Found. Phys.* **2**, (1972), 1-7.
[3] Bleher P.M., Ganikhodjaev N.N. On pure phases of the Ising model on the Bethe lattice. *Theor. Probab. Appl.* **35** (1990), 216-227.
[4] Gandolfo G., Rozikov U.A., Ruiz J. On p-adic Gibbs measures for hard core model on a Cayley tree. *Markov Processes Related Fields.* **18**(4) (2012), 701-720.
[5] Ganikhodjaev N.N., Mukhamedov F.M., Rozikov U.A., Phase Transitions in the Ising Model on Z over the p-adic Number Field, *Uzb. Mat. Zh.*, No. 4, (1998), 23-29.
[6] Gras Georges, Mesures p-adiques. (French) *Théorie des nombres, Année* 1991/1992, 107 pp., Publ. Math.Fac. Sci. Besançon, Univ. Franche-Comté, Besançon.
[7] Georgii H.-O., Gibbs Measures and Phase Transitions (W. de Gruyter, Berlin, 1988).
[8] Khamraev M., Mukhamedov F.M., Rozikov U.A. On the uniqueness of Gibbs measures for $p-$adic non homogeneous $\lambda-$ model on the Cayley tree. *Letters in Math. Phys.* **70** (2004), 17-28.
[9] Khrennikov A. Yu., p-Adic Valued Probability Measures, *Indag. Math.*, New Ser. **7** (1996), 311-330.
[10] Khrennikov A. Yu., p-Adic Valued Distributions in Mathematical Physics (Kluwer, Dordrecht, 1994).
[11] Khrennikov A. Yu., Mukhamedov F.M., Mendes J.F.F., On p-adic Gibbs Measures of the Countable State Potts Model on the Cayley Tree, *Nonlinearity* **20**, (2007), 2923-2937.
[12] Khrennikov A. Yu., Yamada S., van Rooij A., The Measure-Theoretical Approach to p-adic Probability Theory, *Ann. Math. Blaise Pascal* **6**, (1999), 21-32.
[13] Koblitz N., p-Adic Numbers, p-adic Analysis, and Zeta-Functions (Springer, Berlin, 1977).
[14] Mukhamedov F.M. On the Existence of Generalized Gibbs Measures for the One-Dimensional p-adic Countable State Potts Model. *Proc. Steklov Inst. Math.*, **265** (2009), 165-176.
[15] Mukhamedov F.M., Rozikov U.A., On Gibbs Measures of p-adic Potts Model on the Cayley Tree, *Indag. Math.*, New Ser. **15** (2004), 85-100.
[16] Mukhamedov F.M., Rozikov U.A., On Inhomogeneous p-adic Potts Model on a Cayley Tree, *Infin. Dimens. Anal. Quantum Probab. Relat. Top.* **8**, (2005), 277-290.
[17] Mukhamedov F.M., Rozikov U.A., Mendes J.F.F., On Phase Transitions for p-adic Potts Model with Competing Interactions on a Cayley Tree, *in p-Adic Mathematical Physics*: Proc. 2nd Int. Conf., Belgrade, 2005 (Am. Inst. Phys., Melville, NY, 2006), AIP Conf. Proc. 826, pp. 140-150.
[18] Mukhamedov F.M., Saburov M., Khakimov O.N., On p-adic IsingVannimenus model on an arbitrary order Cayley tree. *J. Stat. Mech.*, 2015, P05032.
[19] Mukhamedov F.M., On dynamical systems and phase transitions for $Q+1$-state p-adic Potts model on the Cayley tree. *Math. Phys. Anal. Geom.* **53** (2013), 49–87.
[20] van Rooij A. C. M., Non-Archimedean Functional Analysis (M. Dekker, New York, 1978).
[21] Rozikov U.A. Gibbs measures on Cayley trees. *World Sci. Publ.* Singapore. 2013, 404 pp.
[22] Schikhof W.H., Ultrametric Calculus (Cambridge Univ. Press, Cambridge, 1984).
[23] Shiryaev, A.N. *Probability*, 2nd ed. Graduate Texts in Mathematics, vol. 95. Springer, New York, 1996.
[24] Tugyonov Z. T. On periodic p-adic distributions. *p-Adic Numbers Ultrametric Anal. Appl.* **5**(3) (2013), 218–225.
[25] Tugyonov Z.T. Periodicity of Bernulli's distribution. *Uzbek Math. Jour.* 2 (2013), 107–111.
[26] Tugyonov Z.T. Non Uniqueness of p-adic Gibbs distribution for the Ising model on the lattice Z^d. *Jour. Siber. Federal Univ.* **9**(1) (2016), 123-127.
[27] Vladimirov V.S., Volovich I. V., Zelenov E. V., p-Adic Analysis and Mathematical Physics (Nauka, Moscow, 1994; World Sci., Singapore, 1994).

U. A. Rozikov and Z. T. Tug'yonov, Institute of mathematics, 29, Do'rmon Yo'li str., 100125, Tashkent, Uzbekistan.

E-mail address: rozikovu@yandex.ru zohidboy@mail.ru